



Dating of a young, subarctic peat deposit using the radiocarbon wiggle-match technique

Kokfelt, Ulla; Johansson, Anette; Muscheler, Raimund; Hammarlund, Dan

Published in:
LUCCI Annual Report 2011-2012

Publication date:
2012

Document version
Publisher's PDF, also known as Version of record

Citation for published version (APA):
Kokfelt, U., Johansson, A., Muscheler, R., & Hammarlund, D. (2012). Dating of a young, subarctic peat deposit using the radiocarbon wiggle-match technique. In M. Hellström, M-L. Linderson, & A. Lindroth (Eds.), *LUCCI Annual Report 2011-2012: Lund Centre for Studies of Carbon Cycle and Climate Interactions* (pp. 82-83).
<http://978-91-85793-24-2>

Dating of a young, subarctic peat deposit using the radiocarbon wiggle-match technique

Ulla Kokfelt (a), Anette Johansson (b), Raimund Muscheler (b) and Dan Hammarlund (b)

(a) Department of Geography and Geology, University of Copenhagen, Denmark

(b) Quaternary Sciences, Department of Geology, Lund University

The attribution of forcing factors in paleo-environmental and paleo-climatic studies is often hampered by large uncertainties in the chronological control of proxy records from natural archives. Reducing these dating uncertainties therefore remains a continuous challenge to such studies. One way of reducing dating uncertainties is by wiggle-matching high-resolution ^{14}C measurements against past variations in the atmospheric ^{14}C concentrations [1].

Here, we use ^{14}C wiggle-match dating of closely spaced samples in an attempt to obtain a more solid chronological control of a relatively recent shift in vegetation from a poor *Sphagnum* fen to an ombrotrophic vegetation community separated by a peat layer enriched in diatoms in a peat sequence from Stordalen in northern Sweden ($68^\circ 21' \text{ N}$, $19^\circ 03' \text{ E}$) [2,3]. The major question was whether it is possible to detect even small ^{14}C variations in the calibration curve across significant shifts in vegetation with different trophic status. An age model has previously been constructed based on interpolation between individually calibrated ^{14}C datings below and a ^{210}Pb chronology above the vegetation shift [3].




Figure 1. ^{14}C wiggle-match dating (green dots indicate individual ^{14}C dates) obtained on densely spaced samples from a peat sequence in Stordalen [3], against "wiggles" in the IntCal09 calibration curve prior AD1950 [6] and the NH1 calibration curve after AD1950 [7].

12 contiguous peat samples were collected across the vegetation shift and dated with the Lund University SSAMS system [4]. The results were combined with existing ^{14}C radiocarbon datings [3] and processed in OxCal4.1 [5] using the IntCal09 curve before AD1950 [6] and the post-bomb atmospheric NH1 curve after AD1950 [7].

With the wiggle-match approach, we captured a small but distinct “wiggle” in the ^{14}C calibration curve around AD1600, and we also captured the apparent onset of the bomb-induced increase in atmospheric ^{14}C after 1950 (Fig. 1). Although the results show a promising correspondence with the ^{14}C calibration curve, there are also features that call for further attention. The uppermost five dates showed similar results, which may be a result of contamination by bomb-produced ^{14}C through root penetration. Even minute amounts of roots containing increased amounts of ^{14}C from the bomb tests may have caused these similar dates.

Despite these challenges our results demonstrate that the ^{14}C wiggle-match dating technique is indeed a promising tool for obtaining well-constrained chronologies of recent peat deposits in subarctic Sweden. Future work will focus on improved age modeling of the data by taking advantage of our detailed knowledge of paleo-ecological and biogeochemical changes.

References

- [1] Blaauw M, van Geel B, Mauquoy D, van der Plicht J (2004) Carbon-14 wiggle-match dating of peat deposits: advantages and limitations. *Journal of Quaternary Science* 19 (2): 177-181.
- [2] Kokfelt U, Struyf E and Randsalu L (2009) Diatoms in peat – Dominant producers in a changing environment? *Soil Biology and Biochemistry* 41: 1764-1766.
- [3] Kokfelt U, Reuss N, Struyf E, Sonesson M, Rundgren M, Skog G, Rosén P and Hammarlund D (2010) Wetland development, permafrost history and nutrient cycling inferred from late Holocene peat and lake sediment records in subarctic Sweden. *Journal of Paleolimnology* 44: 327-342.
- [4] Skog G (2007) The single stage AMS machine at Lund University: Status report. *Nuclear Instruments and Methods in Physics Research B* 259: 1-6.
- [5] Bronk Ramsey, C (2009) Bayesian analysis of radiocarbon dates. *Radiocarbon* 51(1): 337-360.
- [6] Reimer P.J., Baillie M.G.L., Bard E, Bayliss A., Beck J.W., Blackwell P.G., Bronk Ramsey C., Buck C.E., Burr G.S., Edwards R.L., Friedrich M., Grootes P.M., Guilderson T.P., Hajdas I., Heaton T.J., Hogg A.G., Hughen K.A., Kaiser K.F., Kromer B., McCormac F.G., Manning S.W., Reimer R.W., Richards D.A., Southon J.R., Talamo S., Turney C.S.M., van der Plicht J., Weyhenmeyer C.E. (2009) IntCal09 and Marine09 radiocarbon age calibration curves, 0-50,000 years Cal BP. *Radiocarbon* 51(4): 1111–1150.
- [7] Hua Q and Barbetti H (2004) Review of tropospheric bomb ^{14}C data for carbon cycle modeling and age calibration purposes. *Radiocarbon* 46(3): 1273–1298.